arXiv:1201.1888v2 [math.RA] 1 Feb 2012

THE SOCLE AND SEMISIMPLICITY OF A KUMJIAN-PASK ALGEBRA

JONATHAN H. BROWN AND ASTRID AN HUEF

Abstract. The Kumjian-Pask algebra $\mathrm{KP}(\Lambda)$ is a graded algebra associated to a higher-rank graph Λ and is a generalization of the Leavitt path algebra of a directed graph. We analyze the minimal left-ideals of $\mathrm{KP}(\Lambda)$, and identify its socle as a graded ideal by describing its generators in terms of a subset of vertices of the graph. We characterize when $\mathrm{KP}(\Lambda)$ is semisimple, and obtain a complete structure theorem for a semisimple Kumjian-Pask algebra.

1. Introduction

The Leavitt path algebra $L(E)$ of a directed graph E over a field is an algebra defined by generators and relations encoded in the graph. Abrams and Aranda Pino [2] constructed Leavitt path algebras in analogy with the graph C^*-algebras introduced in [23]. Leavitt path algebras are so called because they generalize the algebras without invariant basis number studied by Leavitt in [24]. Both Leavitt path algebras and graph C^*-algebras provide a rich source of examples with substantial structure theories and have attracted interest from a broad range of researchers. Results and techniques used to study Leavitt path algebras are now applied to study the C^*-algebras of directed graphs (for example [7, 9]) and vice-versa (for example, [2, 3, 32]). In this paper we study analogues of Leavitt path algebras associated to higher-rank graphs; these algebras are called Kumjian-Pask algebras.

A higher-rank graph Λ is a combinatorial object in which "paths" have a k-dimensional shape or degree, and a 1-graph reduces to a directed graph. Kumjian and Pask introduced k-graphs and the associated C^*-algebras $C^*(\Lambda)$ in [22] to provide models for higher-rank versions of Cuntz-Krieger algebras studied by Robertson and Steger in [28]. The C^*-algebras of k-graphs have been extensively studied (see, for example, [15, 16, 18, 19, 21, 25, 26, 27, 30, 33]), but open problems remain. For example, Evans and Sims have recently found necessary conditions on Λ for $C^*(\Lambda)$ to be approximately finite-dimensional, but the converse seems elusive unless the graph is finitely aligned and has finitely many vertices [18]. Evans and Sims also proved that if Λ is a finitely-aligned k-graph with finitely many vertices, then $C^*(\Lambda)$ is finite-dimensional if and only if Λ contains no "cycles" (and if so then $C^*(\Lambda)$ is isomorphic to a direct sum of matrix algebras indexed by the set of sources of of the graph).

The Kumjian-Pask algebra $\mathrm{KP}(\Lambda)$ is an algebraic version of the C^*-algebra associated to a row-finite k-graph with no sources. Defined and studied in [10], $\mathrm{KP}(\Lambda)$ has a universal property based on a family of generators satisfying suitable Cuntz-Krieger relations, is $\mathbb{Z}^k$-graded, and its two-sided graded ideal structure is given by saturated

---

Date: 9 January, 2012.

2010 *Mathematics Subject Classification.* 16S10, 16W50, 16D70.

Key words and phrases. Kumjian-Pask algebra, Leavitt path algebra, higher-rank graph, k-graph, directed graph, socle, left ideals, semisimplicity.

This research was supported by the University of Otago.

and hereditary subsets of vertices of the k-graph. Example 7.1 in [10] shows that there are Kumjian-Pask algebras that do not arise as Leavitt path algebras.

In this paper we identify the minimal left-ideals of the Kumjian-Pask algebra as those associated to vertices called "line points", use this analysis to identify the socle of the algebra as a graded ideal, and characterize when the algebra is semisimple. We also obtain a complete structure theorem for semisimple Kumjian-Pask algebras reminiscent of the Wedderburn-Artin Theorem.

The socle of a Leavitt path algebra has been studied in [5, 11, 12]. We follow [12] in our analysis of the minimal left-ideals in a Kumjian-Pask algebra and [4] to prove our structure theorem for semisimple algebras. But our methods are often simpler than those of [4] and [12], even when $k = 1$. The proofs in [4] and [12] rely on delicate constructions of specific cycles which we replace with comparison of degrees of paths. Also, our analysis of the direct sum in a semisimple algebra goes a bit further than [4] because we can write our direct sum over an explicit index set of equivalence classes of vertices in Theorem 4.10. In particular, Theorem 4.10 shows first that a finite-dimensional Kumjian-Pask algebra cannot arise from a k-graph with no sources, and second, that a semisimple Kumjian-Pask algebra is isomorphic to a Leavitt path algebra (see Remark 4.11).

2. Preliminaries on k-graphs and Kumjian-Pask algebras

We start by establishing our conventions and proving some basic properties of Kumjian-Pask algebras.

Let k be a positive integer. We consider the additive semigroup $\mathbb{N}^k$ as a category with one object. Following [22, Definition 1.1], we say a countable category $\Lambda = (\Lambda^0, \Lambda, r, s)$, with objects Λ^0, morphisms Λ, range map r and source map s, is a *k-graph* if there exists a functor $d : \Lambda \to \mathbb{N}^k$ with the *unique factorization property*: if $d(\lambda) = m + n$ for some $m, n \in \mathbb{N}^k$, then there exist unique $\mu, \nu \in \Lambda$ such that $r(\nu) = s(\mu)$ and $d(\mu) = m, d(\nu) = n$ with $\lambda = \mu\nu$. Since we think of Λ as a generalized graph, we call $\lambda \in \Lambda$ a *path* in Λ and $v \in \Lambda^0$ a *vertex.*

Let $n \in \mathbb{N}^k$. We use the factorization property to identify Λ^0 and the paths of degree 0, and then define $\Lambda^n := d^{-1}(n)$ and $\Lambda^{\neq n} = \Lambda \setminus \Lambda^n$. For $X, Y \subset \Lambda^0$ we denote by

$$X\Lambda := \{\lambda \in \Lambda : r(\lambda) \in X\}, \quad \Lambda Y := \{\lambda \in \Lambda : s(\lambda) \in Y\}, \quad X\Lambda Y := X\Lambda \cap \Lambda Y,$$

and $X\Lambda^n := X\Lambda \cap \Lambda^n$, $\Lambda^n Y := \Lambda Y \cap \Lambda^n$, and $X\Lambda^n Y := X\Lambda Y \cap \Lambda^n$. For simplicity, we write $v\Lambda$ for $\{v\}\Lambda$.

A k-graph Λ is *row-finite* if $|v\Lambda^n| < \infty$ for all $v \in \Lambda^0, n \in \mathbb{N}^k$ and has *no sources* if $v\Lambda^n \neq \emptyset$ for all $v \in \Lambda^0, n \in \mathbb{N}^k$. We assume throughout that Λ is a row-finite k-graph with no sources, that is $0 < |v\Lambda^n| < \infty$ for all $v \in \Lambda^0, n \in \mathbb{N}^k$.

For $m = (m_1, \ldots, m_k), n = (n_1, \ldots, n_k) \in \mathbb{N}^k$ we say $m \leq n$ if $m_i \leq n_i$ for all i. We denote the join of m and n by $m \vee n$: so $m \vee n = (\max\{m_1, n_1\}, \ldots \max\{m_k, n_k\})$.

Examples 2.1. These examples first appear in [22, Example 1.3 and Examples 1.7]. We discuss them here to establish notation.

(1) Let $E = (E^0, E^1, r_E, s_E)$ be a directed graph. Our convention is that a path μ in E of length $|\mu|$ is a concatenation $\mu = \mu_1\mu_2 \cdots \mu_{|\mu|}$ of edges $\mu_i \in E^1$ with $s_E(\mu_i) = r_E(\mu_{i+1})$. (The usual convention in papers on Leavitt path algebras is to write $\mu = \mu_1\mu_2 \cdots \mu_{|\mu|}$ with $r_E(\mu_i) = s_E(\mu_{i+1})$. We deviate from the usual convention so that concatenation of paths is compatible with composition of morphisms.) Take Λ_E^0 to be the set of vertices, Λ_E the collection of finite

paths, composition to be concatenation of paths, and $r = r_E$, $s = s_E$. Then $\Lambda_E = (E^0, \Lambda, r, s)$ is a category. Equipped with the length function $d : \Lambda_E \to \mathbb{N}$, Λ_E is a 1-graph. Conversely, every 1-graph Λ gives a directed graph E_Λ using the morphisms of degree 1 for the edges.

(2) Let $\Omega_k = \{(m, n) \in \mathbb{N}^k \times \mathbb{N}^k : m \leq n\}$. Then Ω_k is category with objects $\mathbb{N}^k$, morphisms $(m, n) \in \Omega_k$, $r(m, n) = m$ and $s(m, n) = n$, and composition $(m, n)(n, p) = (m, p)$. Equipped with degree map d, defined by $d(m, n) = n - m$, Ω_k is a k-graph.

As in [22, Definition 2.1], define the *infinite path space* of Λ to be

$$\Lambda^\infty = \{x : \Omega_k \to \Lambda : x \quad \text{is a degree-preserving functor}\}$$

and $v\Lambda^\infty = \{x \in \Lambda^\infty : x(0) = v\}$. Since we assume that Λ has no sources, $v\Lambda^\infty \neq \emptyset$ for all $v \in \Lambda^0$. As in [22, Remark 2.2], $x = y \in \Lambda^\infty$ if and only if $x(0, n) = y(0, n)$ for all $n \in \mathbb{N}^k$. For $p \in \mathbb{N}^k$ we define a map $\sigma^p : \Lambda^\infty \to \Lambda^\infty$ by $\sigma^p(x)(m, n) = x(m + p, n + p)$. Note that $x = x(0, p)\sigma^p(x)$. We say that $x \in \Lambda^\infty$ is *periodic* if there exists $p \neq q \in \mathbb{N}^k$ such that $\sigma^p(x) = \sigma^q(x)$: that is for all $m, n \in \mathbb{N}^k$ $x(m + p, n + p) = x(m + q, n + q)$; x is called *aperiodic* otherwise. A k-graph Λ is called *aperiodic* if for every vertex $v \in \Lambda^0$ there exists an aperiodic path in $v\Lambda^\infty$.

Example 2.2. Let Λ be a 1-graph and suppose that $x \in \Lambda^\infty$ is periodic, that is, there exists $p \neq q \in \mathbb{N}$ such that $\sigma^p(x) = \sigma^q(x)$. Since $\mathbb{N}$ is totally ordered, we may assume $p < q$. Then $x(p, q)$ is a cycle in the sense that $r(x(p, q)) = s(x(p, q))$.

Let Λ be a row-finite k-graph with no sources and R a commutative ring with 1. Let $G(\Lambda^{\neq 0}) := \{\lambda^* : \lambda \in \Lambda^{\neq 0}\}$ be a copy of the paths called *ghost paths*. We extend r, s and d to $G(\Lambda^{\neq 0})$ by $r(\lambda^*) = s(\lambda)$, $s(\lambda^*) = r(\lambda)$ and $d(\lambda^*) = -d(\lambda)$. We also extend d to words $w = w_1 \cdots w_{|w|}$ on $\Lambda^0 \cup \Lambda^{\neq 0} \cup G(\Lambda^{\neq 0})$ by $d(w) = \sum_i d(w_i)$.

We now recall [10, Definition 3.1]. A *Kumjian-Pask* Λ*-family* (P, S) in an R-algebra A consists of functions $P : \Lambda^0 \to A$ and $S : \Lambda^{\neq 0} \cup G(\Lambda^{\neq 0}) \to A$ such that

(KP1) $\{P_v : v \in \Lambda^0\}$ is a set of mutually orthogonal idempotents;

(KP2) for $\lambda, \mu \in \Lambda^{\neq 0}$ with $r(\mu) = s(\lambda)$ we have

$$S_\lambda S_\mu = S_{\lambda\mu} \quad S_{\mu^*} S_{\lambda^*} = S_{(\lambda\mu)^*} \quad P_{r(\lambda)} S_\lambda = S_\lambda = S_\lambda P_{s(\lambda)} \quad P_{s(\lambda)} S_{\lambda^*} = S_{\lambda^*} = S_{\lambda^*} P_{r(\lambda)};$$

(KP3) for all $\lambda, \mu \in \Lambda^{\neq 0}$ with $d(\lambda) = d(\mu)$ we have $S_{\lambda^*} S_\mu = \delta_{\lambda,\mu} P_{s(\lambda)}$;

(KP4) for all $v \in \Lambda^0$ and $n \in \mathbb{N}^k \setminus \{0\}$ we have $P_v = \sum_{\lambda \in v\Lambda^n} S_\lambda S_{\lambda^*}$.

It is proved in [10, Theorem 3.4] that there is an R-algebra $\mathrm{KP}_R(\Lambda)$ generated by a Kumjian-Pask Λ-family (p, s) with the following universal property: whenever (Q, T) is a Kumjian-Pask Λ-family in an R-algebra A, there is a unique R-algebra homomorphism $\pi_{Q,T} : \mathrm{KP}_R(\Lambda) \to A$ such that

$$\pi_{Q,T}(p_v) = Q_v, \quad \pi_{Q,T}(s_\lambda) = T_\lambda, \quad \pi_{Q,T}(s_{\mu^*}) = T_{\mu^*}$$

for $v \in \Lambda^0$ and $\lambda, \mu \in \Lambda^{\neq 0}$. The generating family (p, s) is called the universal Kumjian-Pask family. It is a consequence of the *Kumjian-Pask relations* (KP1)–(KP4) that $\mathrm{KP}_R(\Lambda) = \mathrm{span}_R\{s_\mu s_\nu^* : \mu, \nu \in \Lambda \text{ with } s(\mu) = s(\nu)\}$ (with the convention that if $v \in \Lambda_0$ then $s_v = p_v$). Indeed, if $s(\mu) = s(\nu)$ then $s_\mu s_{\nu^*} \neq 0$ in $\mathrm{KP}_R(\Lambda)$. For every nonzero $a \in \mathrm{KP}_R(\Lambda)$ and $n \in \mathbb{N}^k$ there exist $m \geq n$ and a finite subset $F \subset \Lambda \times \Lambda^m$ such that $s(\alpha) = s(\beta)$ for all $(\alpha, \beta) \in F$ and

$$a = \sum_{(\alpha,\beta) \in F \subset \Lambda \times \Lambda^m} r_{\alpha,\beta} s_\alpha s_{\beta^*} \text{ with } r_{\alpha,\beta} \in K \setminus \{0\}; \tag{2.1}$$

in this case we say that a is written in *normal form* [10, Lemma 4.2]. The condition $F \subset \Lambda \times \Lambda^m$ says that all the β appearing in (2.1) are of the same degree m. It is proved in [10, Theorem 3.4] that the subgroups

$$\mathrm{KP}_R(\Lambda)_n := \mathrm{span}_R\{s_\lambda s_{\mu^*} : \lambda, \mu \in \Lambda \text{ and } d(\lambda) - d(\mu) = n\} \qquad (n \in \mathbb{Z}^k)$$

give a $\mathbb{Z}^k$-grading of $\mathrm{KP}_R(\Lambda)$.

Lemma 4.3 of [10] says that for all nonzero $a \in \mathrm{KP}_R(\Lambda)$ there exist $\mu, \nu \in \Lambda$ such that $s_{\mu^*} a s_\nu \neq 0$; the proof of this lemma actually gives a stronger result, and we can then strengthen the conclusion of [10, Proposition 4.9] as well.

Lemma 2.3. *Suppose* $0 \neq a = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta} s_\alpha s_{\beta^*} \in \mathrm{KP}_R(\Lambda)$ *is in normal form.*

1. *For all* $(\mu, \nu) \in F$,
$$s_{\mu^*} a s_\nu = r_{\mu,\nu} p_{s(\mu)} + \sum_{\substack{(\alpha,\nu)\in F \\ d(\alpha)\neq d(\mu)}} r_{\alpha,\nu} s_{\mu^*} s_\alpha \neq 0.$$
2. *If there exists* $v \in \{s(\alpha) : (\alpha, \beta) \in F\}$ *such that* $v\Lambda^\infty$ *contains an aperiodic path, then there exist* $\gamma, \eta \in \Lambda$ *such that* $s_{\gamma^*} a s_\eta = r p_{s(\gamma)} \neq 0$.

Proof. (1) Fix $(\mu, \nu) \in F$. If $(\alpha, \beta) \in F$ then $d(\beta) = d(\nu)$. If $\mu \neq \alpha$ and $s_{\mu^*} s_\alpha \neq 0$ then $d(\mu) \neq d(\alpha)$ by (KP3). So another two applications of (KP3) give

$$s_{\mu^*} a s_\nu = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta} s_{\mu^*} s_\alpha (s_{\beta^*} s_\nu) = \sum_{(\alpha,\nu)\in F} r_{\alpha,\nu} s_{\mu^*} s_\alpha = r_{\mu,\nu} p_{s(\mu)} + \sum_{\substack{(\alpha,\nu)\in F \\ d(\alpha)\neq d(\mu)}} r_{\alpha,\nu} s_{\mu^*} s_\alpha.$$

Therefore the zero-graded component of $s_{\mu^*} a s_\nu$ is precisely $r_{\mu,\nu} p_{s(\mu)}$. Since $r_{\mu,\nu} \neq 0$, $r_{\mu,\nu} p_{s(\mu)} \neq 0$ by [10, Theorem 3.4]. Thus $s_{\mu^*} a s_\nu \neq 0$.

(2) Pick $(\mu, \nu) \in F$ such that $v = s(\mu)$. Then for this $(\mu, \nu) \in F$ we have

$$0 \neq s_{\mu^*} a s_\nu = r_{\mu,\nu} p_{s(\mu)} + \sum_{(\alpha,\nu)\in F} r_{\alpha,\nu} s_{\mu^*} s_\alpha$$

by (1). The result now follows as in the proof of [10, Proposition 4.9] after observing that while [20, Lemma 6.2] is stated for aperiodic k-graphs the proof only requires the existence of an aperiodic path at the vertex v. □

Lemma 2.4. $\mathrm{KP}_R(\Lambda)$ *has a countable set of local units.*

Proof. Let $a \in \mathrm{KP}_R(\Lambda)$, and write a in normal form $a = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta} s_\alpha s_{\beta^*}$. For a finite subset X of Λ^0 define $u_X = \sum_{v\in X} p_v$. Take $W = \{r(\alpha), r(\beta) : (\alpha, \beta) \in F\}$. Then $u_W a = a = a u_W$. Thus $\{u_X : X \text{ is a finite subset of } \Lambda^0\}$ is a set of local units of $\mathrm{KP}_R(\Lambda)$; it is countable because Λ^0 is countable. □

A subset H of Λ^0 is *hereditary* if $r(\lambda) \in H$ for some $\lambda \in \Lambda$ implies $s(\lambda) \in H$. A subset H of Λ^0 is *saturated* if $v \in \Lambda^0, n \in \mathbb{N}^k$ and $s(v\Lambda^n) \subset H$ implies $v \in H$. For every subset $W \subset \Lambda^0$, there exists a smallest subset $\overline{W}$ of Λ^0 such that $\overline{W}$ is both hereditary and saturated; we call $\overline{W}$ the *saturated hereditary closure* of W.

Lemma 2.5. *Let* W *be a subset of* Λ^0 *and let* $\Sigma^0(W)$ *be the smallest hereditary subset of* Λ^0 *containing* W*. If* $v \in \overline{W}$ *then* $v\Lambda\Sigma^0(W) \neq \emptyset$.

Proof. For $N \geq 1$ set $\Sigma^N(W) = \bigcup_{i=1}^k \{v \in \Lambda^0 : s(v\Lambda^{e_i}) \subset \Sigma^{N-1}(W)\}$. Since $\Sigma^0(W)$ is hereditary, $\overline{W} = \bigcup_{N\geq 1} \Sigma^N(W)$ by [26, Lemma 5.1].

Fix $v \in \overline{W}$. Choose N such that $v \in \Sigma^N(W)$. Then there exists $i_N \in \{1, \dots, k\}$ such that $s(v\Lambda^{e_{i_N}}) \subset \Sigma^{N-1}(W)$. In particular, there exists $\mu_N \in v\Lambda\Sigma^{N-1}(W)$. If $N = 1$ we are done. If not, since $s(\mu_N) \in \Sigma^{N-1}(W)$, there exists i_{N-1} such that $s(s(\mu_N)\Lambda^{e_{i_{N-1}}}) \subset \Sigma^{N-1}(W)$. In particular, there exists $\mu_{N-1} \in s(\mu_N)\Lambda\Sigma^{N-2}(W)$. Now $\mu_N\mu_{N-1} \in v\Lambda\Sigma^{N-2}(W)$. Continue to obtain $\mu := \mu_N\mu_{N-1}\dots\mu_1 \in v\Lambda\Sigma^0(W)$. □

For a saturated hereditary subset $H \subset \Lambda^0$ define I_H to be the ideal of $\mathrm{KP}_R(\Lambda)$ generated by $\{p_v : v \in H\}$. For an ideal I in $\mathrm{KP}_R(\Lambda)$ define $H_I = \{v : p_v \in I\}$. Then H_I is saturated hereditary by [10, Lemma 5.2], and if H is saturated hereditary then $H_{I_H} = H$ by [10, Lemma 5.4]. An ideal I in $\mathrm{KP}_R(\Lambda)$ is *graded* if $I = \oplus_{n\in\mathbb{Z}^k} I \cap \mathrm{KP}_R(\Lambda)_n$, which occurs if and only if I is generated by homogeneous elements. An ideal I is *basic* if $tp_v \in I$ and $t \in R \setminus \{0\}$ implies that $p_v \in I$. If $R = K$ is a field then every ideal in $\mathrm{KP}_K(\Lambda)$ is basic, and hence [10, Theorem 5.1] implies that the map $H \mapsto I_H$ is a lattice isomorphism of the saturated hereditary subsets of Λ^0 onto the graded ideals of $\mathrm{KP}_K(\Lambda)$. Here we restrict our attention to Kumjian-Pask algebras over a field K.

Lemma 2.6. *Let K be a field. Let $W \subset \Lambda^0$ and let J be the ideal generated by $\{p_v : v \in W\}$. Then J is a graded ideal of $\mathrm{KP}_K(\Lambda)$, and $J = I_{\overline{W}}$ and $H_J = \overline{W}$.*

Proof. Suppose $W \subset \Lambda^0$ and J is the ideal generated by $\{p_v : v \in W\}$. We have $W \subset H_J = \{v : p_v \in J\}$, and H_J is saturated hereditary. Thus $\overline{W} \subset H_J$. Since $\{p_v : v \in W\}$ is a set of homogeneous elements the ideal J generated by it is graded. By [10, Theorem 5.1], $H \mapsto I_H$ is a lattice isomorphism of the saturated hereditary subsets of Λ^0 onto the graded ideals of $\mathrm{KP}_K(\Lambda)$, and $J = I_{H_J}$. Now $J \subset I_{\overline{W}} \subset I_{H_J} = J$. Thus $J = I_{\overline{W}}$ and $H_J = \overline{W}$. □

A ring R is *nondegenerate* if $aRa = \{0\}$ implies that $a = 0$. A ring R is *semiprime* if whenever I is an ideal in R such that $I^2 = \{0\}$, then $I = \{0\}$.

Lemma 2.7. *Let K be a field. Then $\mathrm{KP}_K(\Lambda)$ is nondegenerate and semiprime.*

Proof. A ring with local units is nondegenerate if and only if it is semiprime by, for example, [14, Proposition 6.2.20]. Since $\mathrm{KP}_K(\Lambda)$ has local units by Lemma 2.4, it suffices to show that $\mathrm{KP}_K(\Lambda)$ is nondegenerate.

First suppose that $a\,\mathrm{KP}_K(\Lambda)a = \{0\}$ where $a \in \mathrm{KP}_K(\Lambda)_n$ is homogeneous of degree $n \in \mathbb{Z}^k$. By way of contradiction, suppose $a \neq 0$ and write $a = \sum_{(\alpha,\beta)\in F} r_{(\alpha,\beta)} s_\alpha s_{\beta^*}$ in normal form. Since $a \in \mathrm{KP}_K(\Lambda)_n$, all of the summands of a are in $\mathrm{KP}_K(\Lambda)_n$ as well. Since the β all have the same degree, so do the α. Fix $(\mu, \nu) \in F$. Then $s_{\mu^*} a s_\nu$ is homogeneous, and it follows from Lemma 2.3(1) that $s_{\mu^*} a s_\nu = r_{\mu,\nu} p_{s(\mu)}$. So $r_{\mu,\nu}p_{s(\mu)} \mathrm{KP}_K(\Lambda) r_{\mu,\nu}p_{s(\mu)} = s_{\mu^*} a s_\nu \mathrm{KP}_K(\Lambda) s_{\mu^*} a s_\nu \subset s_{\mu^*} (a\,\mathrm{KP}_K(\Lambda)a)\, s_\nu = \{0\}$. But now

$$r_{\mu,\nu}^2 p_{s(\mu)} = (r_{\mu,\nu}p_{s(\mu)})p_{s(\mu)}(r_{\mu,\nu}p_{s(\mu)}) \in r_{\mu,\nu}p_{s(\mu)} \mathrm{KP}_K(\Lambda) r_{\mu,\nu}p_{s(\mu)} = \{0\}$$

implies that $r_{\mu,\nu}^2 = 0$. Since K is a field, $r_{\mu,\nu} = 0$. But $(\mu, \nu) \in F$ was arbitrary, so $a = 0$, a contradiction. Thus $a\,\mathrm{KP}_K(\Lambda)a = \{0\}$ and a homogeneous implies that $a = 0$.

Next, suppose $a\,\mathrm{KP}_K(\Lambda)a = \{0\}$ for some $a \in \mathrm{KP}_K(\Lambda)$. By way of contradiction again, suppose $a \neq 0$. Decompose a into nonzero homogeneous components $a = \sum_{n\in G} a_n$ with $a_n \in \mathrm{KP}_K(\Lambda)_n$ for some finite subset G of $\mathbb{Z}^k$. If $b \in \mathrm{KP}_K(\Lambda)_m$ for some m, then $aba \in a\,\mathrm{KP}_K(\Lambda)a = \{0\}$. Let $n, n' \in G$ such that $n \neq n'$. Then the degrees of $a_n b a_n$ and $a_{n'} b a_{n'}$ are different. Therefore $a_n b a_n$ is the unique homogeneous component of aba of degree $2d(a_n) + m$, and hence must be zero. Thus $a_n b a_n = 0$ for all homogeneous elements $b \in \mathrm{KP}_K(\Lambda)$. Since $\mathrm{KP}_K(\Lambda)$ is generated by homogeneous elements we have $a_n \operatorname{KP}_K(\Lambda) a_n = \{0\}$. Therefore $a_n = 0$ by the previous paragraph, a contradiction. Hence $\operatorname{KP}_K(\Lambda)$ is nondegenerate. $\square$

3. Minimal left-ideals and the socle of a Kumjian-Pask algebra

Throughout this section Λ is a row-finite k-graph with no sources and K is a field. In analogy with the analysis of the minimal left-ideals of the Leavitt path algebras in [12], we show that minimal left-ideals of the Kumjian-Pask algebra are isomorphic to $\operatorname{KP}_K(\Lambda)p_v$ where v is a vertex called a "line point" in Definition 3.2 below. We then identify the socle of $\operatorname{KP}_K(\Lambda)$ as the graded ideal generated by the vertex idempotents p_v where v is in the saturated hereditary closure of the line points.

Lemma 3.1. *Let $a \in \operatorname{KP}_K(\Lambda)$, and suppose that $\operatorname{KP}_K(\Lambda)a$ is a minimal left-ideal. Then there exist a vertex $v \in \Lambda^0$ and $b \in p_v \operatorname{KP}_K(\Lambda) p_v$ such that $\operatorname{KP}_K(\Lambda)a$ and $\operatorname{KP}_K(\Lambda)b$ are isomorphic as left modules.*

Proof. Suppose that $\operatorname{KP}_K(\Lambda)a$ is a minimal left-ideal. Then $a \neq 0$ and we write $a = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta} s_\alpha s_{\beta^*}$ in normal form. Pick $(\mu,\nu) \in F$ and let $v = s(\mu) = s(\nu)$. Then, by Lemma 2.3(1), $0 \neq s_{\mu^*} a s_\nu \in p_v \operatorname{KP}_K(\Lambda) p_v$. Since $\operatorname{KP}_K(\Lambda)a$ is minimal and $s_{\mu^*}a \neq 0$, $\operatorname{KP}_K(\Lambda)a = \operatorname{KP}_K(\Lambda) s_{\mu^*} a$. Let $b = s_{\mu^*} a s_\nu$, then $c s_{\mu^*} a \mapsto cb$ defines a left-module homomorphism, $\Phi : \operatorname{KP}_K(\Lambda)a \to \operatorname{KP}_K(\Lambda)b$. Indeed, if $c s_{\mu^*} a = d s_{\mu^*} a$ then $(c-d) s_{\mu^*} a = 0$ and hence $cb - db = (c-d) s_{\mu^*} a s_\nu = 0$, so Φ is well-defined. Now Φ is a surjective module homomorphism, and it is injective since the kernel of Φ is a left ideal in the minimal left-ideal $\operatorname{KP}_K(\Lambda)a$. $\square$

By Lemma 3.1 we may assume that minimal left-ideals are of the form $\operatorname{KP}_K(\Lambda)b$ for some $b \in p_v \operatorname{KP}_K(\Lambda) p_v$ and $v \in \Lambda^0$; next we analyze the properties of such v.

Definition 3.2. Suppose that Λ is a k-graph. A vertex $v \in \Lambda^0$ is a *line point* if $v\Lambda^\infty = \{x\}$ and x is aperiodic. We denote the set of line points of Λ by $P_l(\Lambda)$.

The following observations are needed below. In particular, they help in Remark 3.4 where we reconcile Definition 3.2 with the line points of directed graphs as defined in [12, Definition 2.1].

Remark 3.3. As pointed out in [22, Remark 2.2], the values of $x(0,m)$ for $m \in \mathbb{N}^k$ completely determine $x \in \Lambda^\infty$. Thus:

(1) $|v\Lambda^\infty| = 1$ if and only if $|v\Lambda^m| = 1$ for all $m \in \mathbb{N}^k$.
(2) Suppose that $v\Lambda^\infty = \{x\}$. Then $x(m)\Lambda^\infty = \{\sigma^m(x)\}$ for all $m \in \mathbb{N}^k$. Thus, if $x(p) = x(q)$ for some $p \neq q \in \mathbb{N}^k$ then $\sigma^p(x) = \sigma^q(x)$, that is, x is periodic.
(3) The set $P_l(\Lambda)$ is hereditary. To see this, let $\mu \in v\Lambda$ for some line point v. Then $v\Lambda^\infty = \{x\}$ where x is aperiodic. By (2), $s(\mu)\Lambda^\infty = \{\sigma^{d(\mu)}(x)\}$, and $\sigma^{d(\mu)}(x)$ is aperiodic because x is. Thus $s(\mu)$ is a line point, and $P_l(\Lambda)$ is hereditary.
(4) If $|v\Lambda^\infty| = 1$ and $\alpha \in v\Lambda$, then from (1), $v\Lambda^{d(\alpha)} = \{\alpha\}$. So (KP4) implies that
$$p_v = \sum_{\lambda \in v\Lambda^{d(\alpha)}} s_\lambda s_{\lambda^*} = s_\alpha s_{\alpha^*}.$$

Remark 3.4. If Λ is a 1-graph then Definition 3.2 reduces to [12, Definition 2.1]. To see this, let $T(v)$ be the smallest hereditary subset of Λ^0 containing $v \in \Lambda^0$. Notice that $|v\Lambda^\infty| > 1$ if and only if there exists $u \in T(v)$ with $|u\Lambda^m| > 1$ for some $m \in \mathbb{N}$. Now let v be a line point, that is, $v\Lambda^\infty = \{x\}$ where x is aperiodic. Since $|v\Lambda^\infty| = 1$ there can be no "bifurcations" at any $u \in T(v)$, and since x is aperiodic there can be no cycles at any $u \in T(v)$. Thus v is a line point in the sense of [12, Definition 2.1].

Let $a \in p_v \operatorname{KP}_K(\Lambda) p_v$ and suppose that $\operatorname{KP}_K(\Lambda)a$ is a minimal left-ideal. Our next goal is to show that v is a line point, and we do this in two steps: the next proposition shows that $v\Lambda^\infty$ is a singleton $\{y\}$ and Proposition 3.7 shows that y must be aperiodic. Our analysis of the minimal left-ideals culminates in Theorem 3.9.

Proposition 3.5. *Let $v \in \Lambda^0$ and $a \in p_v \operatorname{KP}_K(\Lambda) p_v$.*

(1) Let $m \in \mathbb{N}^k$. Then

$$\Psi : \operatorname{KP}_K(\Lambda)a \to \bigoplus_{\lambda \in v\Lambda^m} \operatorname{KP}_K(\Lambda) s_\lambda s_{\lambda^*} a, \quad \Psi(ca) = (c s_\lambda s_{\lambda^*} a)_{\lambda \in v\Lambda^m}$$

is an isomorphism of left-modules.

(2) If $\operatorname{KP}_K(\Lambda)a$ is a minimal left-ideal then $|v\Lambda^\infty| = 1$.

Proof. (1) Since Ψ is a left-module homomorphism it suffices to write down an inverse for it. Define $\Phi : (c_\lambda s_\lambda s_{\lambda^*} a)_{\lambda \in v\Lambda^m} \mapsto \sum_{\lambda \in v\Lambda^m} c_\lambda s_\lambda s_{\lambda^*} a$. Then $\Phi \circ \Psi(ca) = \Phi((c s_\lambda s_{\lambda^*} a)) = \sum_{\lambda \in v\Lambda^m} c s_\lambda s_{\lambda^*} a = c p_v a = ca$ using (KP4). Also,

$$\begin{aligned}\Psi \circ \Phi((c_\lambda s_\lambda s_{\lambda^*} a)_{\lambda \in v\Lambda^m}) &= \Psi\Big(\sum_{\lambda \in v\Lambda^m} c_\lambda s_\lambda s_{\lambda^*} a\Big) = \Psi\Big(\Big(\sum_{\lambda \in v\Lambda^m} c_\lambda s_\lambda s_{\lambda^*}\Big)a\Big) \\ &= \Big(\sum_{\lambda \in v\Lambda^m} c_\lambda s_\lambda s_{\lambda^*} s_\mu s_{\mu^*} a\Big)_{\mu \in v\Lambda^m} = (c_\lambda s_\lambda s_{\lambda^*} a)_{\lambda \in v\Lambda^m}\end{aligned}$$

using (KP3). Thus Φ is an inverse for Ψ, and Ψ is an isomorphism.

(2) Suppose that $|v\Lambda^\infty| > 1$. Then there exists $m \in \mathbb{N}^k$ such that $|v\Lambda^m| > 1$ (see Remark 3.3(1)). Then $\operatorname{KP}_K(\Lambda)a$ and $\bigoplus_{\lambda \in v\Lambda^m} \operatorname{KP}_K(\Lambda) s_\lambda s_{\lambda^*} a$ are isomorphic by (1). But the direct sum has at least two summands, and hence $\operatorname{KP}_K(\Lambda)a$ is not minimal. □

The next lemma is needed again in §4.

Lemma 3.6. *Let $v \in \Lambda^0$ such that $|v\Lambda^\infty| = 1$ and $N \in \mathbb{N}$. Suppose $\mu_i, \nu_i \in v\Lambda$ such that $s(\mu_i) = s(\nu_i)$ for $i = 1, \dots, N$. Then $s_{\mu_1} s_{\nu_1^*} \cdots s_{\mu_N} s_{\nu_N^*} \neq 0$.*

Proof. We proceed by induction on N. Since $s(\mu_1) = s(\nu_1)$, $0 \neq p_{s(\mu_1)} = s_{\mu_1^*} s_{\mu_1} s_{\nu_1^*} s_{\nu_1}$. In particular $s_{\mu_1} s_{\nu_1^*} \neq 0$, and the lemma holds for $N = 1$.

Let $N \geq 2$, and suppose every set $\{\alpha_i, \beta_i \in v\Lambda : s(\alpha_i) = s(\beta_i), i = 1, \dots N-1\}$ has the property that $s_{\alpha_1} s_{\beta_1^*} \cdots s_{\alpha_{N-1}} s_{\beta_{N-1}^*} \neq 0$. Pick $\{\mu_i, \nu_i \in v\Lambda : s(\mu_i) = s(\nu_i), i = 1, \dots, N\}$ and consider $s_{\mu_1} s_{\nu_1^*} \cdots s_{\mu_N} s_{\nu_N^*}$. Since $r(\nu_1) = r(\mu_2) = v$ there exist $\gamma, \eta \in \Lambda$ such that $\nu_1 \gamma = \mu_2 \eta \in v\Lambda^{d(\nu_1) \vee d(\mu_2)}$. Since $|v\Lambda^\infty| = 1$ we have $s(\nu_1)\Lambda^{d(\gamma)} = \{\gamma\}$ and $s(\mu_2)\Lambda^{d(\eta)} = \{\eta\}$ (see Remark 3.3(1)). Thus

$$\begin{aligned} s_{\mu_1} s_{\nu_1^*} s_{\mu_2} s_{\nu_2^*} \cdots s_{\mu_N} s_{\nu_N^*} &= s_{\mu_1} p_{s(\nu_1)} s_{\nu_1^*} s_{\mu_2} p_{s(\mu_2)} s_{\nu_2^*} \cdots s_{\mu_N} s_{\nu_N^*} \quad \text{(using (KP2))} \\ &= s_{\mu_1} (s_\gamma s_{\gamma^*}) s_{\nu_1^*} s_{\mu_2} (s_\eta s_{\eta^*}) s_{\nu_2^*} \cdots s_{\mu_N} s_{\nu_N^*} \quad \text{(using (KP4) and Remark 3.3(4))} \\ &= s_{\mu_1 \gamma} (s_{(\nu_1 \gamma)^*} s_{\mu_2 \eta}) s_{(\nu_2 \eta)^*} \cdots s_{\mu_N} s_{\nu_N^*} \\ &= s_{\mu_1 \gamma} s_{(\nu_2 \eta)^*} \cdots s_{\mu_N} s_{\nu_N^*} \quad \text{(by (KP3) since } \nu_1 \gamma = \mu_2 \eta.) \end{aligned} \tag{3.1}$$

But (3.1) is nonzero by our inductive hypothesis, and the lemma follows. □

Proposition 3.7. *Let $v \in \Lambda^0$ such that $v\Lambda^\infty = \{y\}$ and let $a \in p_v \operatorname{KP}_K(\Lambda) p_v$. If $\operatorname{KP}_K(\Lambda)a$ is a minimal left-ideal then y is aperiodic.*

Proof. Suppose that y is periodic. Then there exists $n \neq m \in \mathbb{N}^k$ with $\sigma^m(y) = \sigma^n(y)$: in particular, $y(m) = y(n)$. Let $\pi_h : \mathbb{Z}^k \to \mathbb{Z}$ be the projection onto the h-th factor. By possibly interchanging the roles of m and n we can assume that $\pi_l(m - n) > 0$ for some $l \in \{1, \dots, k\}$. Let $\mu := y(0, m)$ and $\nu := y(0, n)$. Since $s(\mu) = s(\nu)$, $s_\mu s_{\nu^*} \neq 0$. Consider $s_\mu s_{\nu^*} a + a \in \mathrm{KP}_K(\Lambda)a$. We have $\{0\} \neq \mathrm{KP}_K(\Lambda)(s_\mu s_{\nu^*} a + a) \subset \mathrm{KP}_K(\Lambda)a$. We will show that this inclusion is proper by showing that $a \notin \mathrm{KP}_K(\Lambda)(s_\mu s_{\nu^*} a + a)$.

Let $a = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta} s_\alpha s_{\beta^*}$ be in normal form. Then for all $(\alpha, \beta), (\mu, \nu) \in F$ we have $d(\beta) = d(\nu)$, and, since $|v\Lambda^\infty| = 1$, we have $\beta = \nu$. So we can assume that

$$a = \sum_{j=1}^{N} r_j s_{\alpha_j} s_{\beta^*} \quad r_j \in K \setminus \{0\}$$

for some fixed β. We may assume that the α_j are ordered so that $\pi_l(d(\alpha_j)) \leq \pi_l(d(\alpha_{j'}))$ for all $1 \leq j \leq j' \leq N$.

By way of contradiction, suppose that $a \in \mathrm{KP}_K(\Lambda)(s_\mu s_{\nu^*} a + a)$, that is, suppose that there exists $b \in \mathrm{KP}_K(\Lambda)$ such that

$$b(s_\mu s_{\nu^*} a + a) = a. \tag{3.2}$$

Then $p_v b p_v (s_\mu s_{\nu^*} a + a) = p_v a = a$ so we may replace b by $p_v b p_v$, that is, we may assume $b \in p_v \mathrm{KP}_K(\Lambda) p_v$. As above there is a $\tau \in \Lambda$ such that b has normal form

$$b = \sum_{i=1}^{M} t_i s_{\gamma_i} s_{\tau^*} \quad \text{with } \pi_l(d(\gamma_i)) \leq \pi_l(d(\gamma_{i'})) \text{ for } 1 \leq i \leq i' \leq M.$$

Our strategy is to compare degrees of the terms on the left and right side of equation (3.2); in particular we will compare the terms whose degree has maximal and minimal l-th component. The terms of $b(s_\mu s_{\nu^*} a + a)$ are either of the form $s_{\gamma_i} s_{\tau^*} s_{\alpha_j} s_{\beta^*}$ (the summands of ba) or $s_{\gamma_i} s_{\tau^*} s_\mu s_{\nu^*} s_{\alpha_j} s_{\beta^*}$ (the summands of $bs_\mu s_{\nu^*} a$). Note that both $s_{\gamma_i} s_{\tau^*} s_{\alpha_j} s_{\beta^*}$ and $s_{\gamma_i} s_{\tau^*} s_\mu s_{\nu^*} s_{\alpha_j} s_{\beta^*}$ are nonzero by Lemma 3.6.

For a homogeneous element $c \in \mathrm{KP}_K(\Lambda)_n$ of degree n we will call $\pi_l(d(c))$ the π_l-degree of c. Since $d(\mu) = m, d(\nu) = n$ and $\pi_l(m - n) > 0$, the terms of $b(s_\mu s_{\nu^*} a + a)$ who have maximal π_l-degree come from $bs_\mu s_{\nu^*} a$. Consider the term of maximal π_l-degree, $r_N t_M s_{\gamma_M} s_{\tau^*} s_\mu s_{\nu^*} s_{\alpha_N} s_{\beta^*}$. Then

$$\pi_l(d(\gamma_M \tau^* \mu \nu^* \alpha_N \beta^*)) = \pi_l(d(\gamma_M \tau^*)) + \pi_l(m - n) + \pi_l(d(\alpha_N \beta^*)).$$

By assumption on the ordering of $\sum r_i s_{\alpha_i} s_{\beta^*}$, the maximal π_l-degree of terms on the right-hand side of equation (3.2) is $\pi_l(d(\alpha_N \beta^*))$. So for (3.2) to hold we must have $\pi_l(d(\gamma_M \tau^*)) + \pi_l(m - n) + \pi_l(d(\alpha_N \beta^*)) = \pi_l(d(\alpha_N \beta^*))$. Hence

$$\pi_l(d(\gamma_M \tau^*)) = -\pi_l(m - n) < 0.$$

But $\pi_l(d(\gamma_M \tau^*))$ is maximal among the π_l-degrees of b, so $\pi_l(d(\gamma_i \tau^*)) < 0$ for all i.

Since $\pi_l(m - n) > 0$, the terms with minimal π_l-degree in $b(s_\mu s_{\nu^*} a + a)$ come from ba. Consider the term of minimal π_l-degree, $r_1 t_1 s_{\gamma_1} s_{\tau^*} s_{\alpha_1} s_{\beta^*}$. From above, $\pi_l(d(\gamma_1 \tau^*)) < 0$, so $\pi_l(d(\gamma_1 \tau^* \alpha_1 \beta^*)) = \pi_l(d(\gamma_1 \tau^*)) + \pi_l(d(\alpha_1 \beta^*)) < \pi_l(d(\alpha_1 \beta^*))$. But $s_{\alpha_1} s_{\beta^*}$ is the term on the right-hand side of (3.2) of minimal π_l-degree. So there is no component of a that has degree $d(\gamma_1 \tau^* \alpha_1 \beta^*)$ and hence (3.2) cannot hold. So there does not exist $b \in \mathrm{KP}_K(\Lambda)$ such that $b(s_\mu s_{\nu^*} a + a) = a$. Hence $\{0\} \neq \mathrm{KP}_K(\Lambda)(a s_\mu s_{\nu^*} + a) \subsetneq \mathrm{KP}_K(\Lambda)a$; in particular, $\mathrm{KP}_K(\Lambda)a$ is not a minimal left-ideal. □

Lemma 3.8. *If* $v \in P_l(\Lambda)$ *then* $p_v \mathrm{KP}_K(\Lambda) p_v = K p_v$.

Proof. Let $v \in P_l(\Lambda)$, that is, $v\Lambda^\infty = \{x\}$ where x is aperiodic. To see $p_v \mathrm{KP}_K(\Lambda) p_v \subset K p_v$, let $p_v s_\alpha s_{\beta^*} p_v$ be nonzero. Then $s(\alpha) = s(\beta)$ and $\alpha, \beta \in v\Lambda$. It follows from Remark 3.3(2) that $\alpha = \beta$. Now $p_v s_\alpha s_{\beta^*} p_v = p_v s_\alpha s_{\alpha^*} p_v = p_v$ by (KP4) and Remark 3.3(4).

Since elements of the form $s_\alpha s_{\beta^*}$ generate $\mathrm{KP}_K(\Lambda)$ we get $p_v \mathrm{KP}_K(\Lambda)p_v \subset Kp_v$. The other inclusion is immediate because $p_v = p_v^3$. $\square$

Theorem 3.9. *Let Λ be a row-finite k-graph with no sources and let K be a field.*

1. *Let $v \in P_l(\Lambda)$. Then $\mathrm{KP}_K(\Lambda)p_v$ is a minimal left-ideal if and only if $v \in P_l(\Lambda)$.*
2. *Let $a \in \mathrm{KP}_K(\Lambda)$. Then $\mathrm{KP}_K(\Lambda)a$ is a minimal left-ideal if and only if there exists $v \in P_l(\Lambda)$ such that $\mathrm{KP}_K(\Lambda)p_v$ is isomorphic to $\mathrm{KP}_K(\Lambda)a$ as left $\mathrm{KP}_K(\Lambda)$-modules.*

Proof. (1) Let $v \in P_l(\Lambda)$. We will show that $\mathrm{KP}_K(\Lambda)ap_v = \mathrm{KP}_K(\Lambda)p_v$ for all nonzero $a \in \mathrm{KP}_K(\Lambda)$; it then follows that $\mathrm{KP}_K(p_v)$ is a minimal left-ideal. Fix $0 \neq a \in \mathrm{KP}_K(\Lambda)$. It is enough to show $p_v \in \mathrm{KP}_K(\Lambda)ap_v$. By Lemma 2.7, $ap_v \mathrm{KP}_K(\Lambda)ap_v \neq \{0\}$. Thus there exists $b \in \mathrm{KP}_K(\Lambda)$ such that $p_v bap_v \neq 0$. By Lemma 3.8, $p_v \mathrm{KP}_K(\Lambda)p_v = Kp_v$ so there exists $t \in K \setminus \{0\}$ such that $tp_v = p_v bap_v$. Now $p_v = (t^{-1}p_v b)ap_v \in \mathrm{KP}_K(\Lambda)ap_v$, giving $\mathrm{KP}_K(\Lambda)ap_v = \mathrm{KP}_K(\Lambda)p_v$. Thus $\mathrm{KP}_K(\Lambda)p_v$ is minimal.

Conversely, suppose that $\mathrm{KP}_K(\Lambda)p_v$ is minimal for some $v \in \Lambda^0$. By item (2) of Proposition 3.5 we have $v\Lambda^\infty = \{y\}$ for some y. Now Proposition 3.7 implies that y is aperiodic. Thus $v \in P_l(\Lambda)$.

(2) Suppose $\mathrm{KP}_K(\Lambda)a$ is minimal. By Lemma 3.1, we can assume $a \in p_w \mathrm{KP}_K(\Lambda)p_w$ for some $w \in \Lambda^0$. Since $\mathrm{KP}_K(\Lambda)a$ is minimal, $w\Lambda^\infty = \{x\}$ by Proposition 3.5(2), and then x is aperiodic by Proposition 3.7. Thus $w \in P_l(\Lambda)$. Since $x \in w\Lambda^\infty$ is aperiodic, Lemma 2.3(2) shows there exist $\gamma, \eta \in \Lambda$ such that $s_{\gamma^*}as_\eta = rp_{s(\gamma)} \neq 0$ for some $r \in K$. But $a \in p_w \mathrm{KP}_K(\Lambda)p_w$ so $\gamma, \eta \in w\Lambda$. Since $\mathrm{KP}_K(\Lambda)a$ is minimal, $\mathrm{KP}_K(\Lambda)a = \mathrm{KP}_K(\Lambda)r^{-1}s_{\gamma^*}a$.

We claim that the map $cs_{\gamma^*}a \mapsto cs_{\gamma^*}as_\eta$ defines an isomorphism from $\mathrm{KP}_K(\Lambda)r^{-1}s_{\gamma^*}a$ to $\mathrm{KP}_K(\Lambda)s_{\gamma^*}as_\eta$. This map is surjective by definition, and is injective because the kernel is a left-ideal in the minimal left-ideal $\mathrm{KP}_K(\Lambda)s_{\gamma^*}a$, hence must be $\{0\}$. Thus $\mathrm{KP}_K(\Lambda)a = \mathrm{KP}_K(\Lambda)r^{-1}s_{\gamma^*}a$ is isomorphic to $\mathrm{KP}_K(\Lambda)s_{\gamma^*}as_\eta = \mathrm{KP}_K(\Lambda)p_{s(\gamma)}$ as claimed.

Finally, $r(\gamma) = w \in P_l(\Lambda)$ and $P_l(\Lambda)$ is hereditary by Remark 3.3(4), and hence $s(\gamma) \in P_l(\Lambda)$. Thus $\mathrm{KP}_K(\Lambda)a$ is isomorphic to $\mathrm{KP}_K(\Lambda)p_{s(\gamma)}$ and $s(\gamma) \in P_l(\Lambda)$. The converse is immediate from (1). $\square$

The *left socle* $\mathrm{Soc}_l(A)$ of an algebra A is the sum of all of its minimal left-ideals; if the set of minimal left-ideals is empty then the left socle is zero. When A is semiprime, the left socle and the right socle coincide [13, Corollary 4.3.4], and then the *socle* $\mathrm{Soc}(A) := \mathrm{Soc}_l(A)$ of A is a two-sided ideal.

By Lemma 2.7, the Kumjian-Pask algebra $\mathrm{KP}_K(\Lambda)$ is semiprime. By Theorem 3.9, we have $\sum_{v\in P_l(\Lambda)} \mathrm{KP}_K(\Lambda)p_v \subset \mathrm{Soc}(\mathrm{KP}_K(\Lambda))$ where $P_l(\Lambda)$ is the set of line points. The example in the proof of [12, Proposition 4.1] shows the reverse containment does not hold in general. In Theorem 3.10 we show that $\mathrm{Soc}(\mathrm{KP}_K(\Lambda))$ is the ideal corresponding to the saturated hereditary closure of the line points.

When $k = 1$ the first item of Theorem 3.10 below is [12, Theorem 4.2] and the second item is [29, Theorem 4.3]. An ideal I in an algebra is *essential* if $a \in A$ and $aI = \{0\}$ implies $a = 0$.

Theorem 3.10. *Let Λ be a row-finite k-graph with no sources and let K be a field.*

1. *Let J be the ideal generated by $\{p_v : v \in P_l(\Lambda)\}$. Then*
$$\mathrm{Soc}(\mathrm{KP}_K(\Lambda)) = J = I_{\overline{P_l(\Lambda)}}.$$
2. *$\mathrm{Soc}(\mathrm{KP}_K(\Lambda))$ is an essential ideal of $\mathrm{KP}_K(\Lambda)$ if and only if every vertex connects to a line point.*

Proof. (1) The proof is very similar to that of [12, Theorem 4.2]. By Lemma 2.6, $I_{\overline{P_l(\Lambda)}} = J$. To see that $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) \subset J$, let L be a minimal left-ideal of $\mathrm{KP}_K(\Lambda)$; we will show that $L \subset J$. Let $a \in L \setminus \{0\}$. Then $\mathrm{KP}_K(\Lambda)a \subset L$. By the nondegeneracy of $\mathrm{KP}_K(\Lambda)$ (Lemma 2.7) we have $\mathrm{KP}_K(\Lambda)a \neq 0$. Thus $L = \mathrm{KP}_K(\Lambda)a$ by the minimality of L. By Theorem 3.9, $L = \mathrm{KP}_K(\Lambda)a$ is isomorphic to $\mathrm{KP}_K(\Lambda)p_v$ for some line point v. Let $\phi : \mathrm{KP}_K(\Lambda)a \to \mathrm{KP}_K(\Lambda)p_v$ be an isomorphism. Then $\phi(a) = bp_v$ for some $b \in \mathrm{KP}_K(\Lambda)$. Note that since v is a line point, $p_v \in J$ and so $a = \phi^{-1}(\phi(a)) = \phi^{-1}(bp_v) = \phi^{-1}(bp_vp_v) = bp_v\phi^{-1}(p_v) \in J$. Thus $L \subset J$ and hence $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) \subset J$.

For the reverse containment, if $p_v \in P_l(\Lambda)$ then $\mathrm{KP}_K(\Lambda)p_v \subset \operatorname{Soc}(\mathrm{KP}_K(\Lambda))$. Thus $p_v = p_vp_v \in \operatorname{Soc}(\mathrm{KP}_K(\Lambda))$ for all $v \in P_l(\Lambda)$. Therefore $\operatorname{Soc}(\mathrm{KP}_K(\Lambda))$ is an ideal containing $\{p_v : v \in P_l(\Lambda)\}$ and so it contains J.

(2) First suppose that every vertex in $\mathrm{KP}_K(\Lambda)$ connects to a line point. Let $a \in \mathrm{KP}_K(\Lambda) \setminus \{0\}$; we will show that $a \operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = aI_{\overline{P_l(\Lambda)}} \neq \{0\}$. First we claim that Λ is aperiodic. To see this, let $u \in \Lambda^0$. Then there exists a path $\lambda \in u\Lambda v$ where v is a line point. But $v\Lambda^\infty = \{x\}$ with x aperiodic, and now λx is an aperiodic path in $u\Lambda^\infty$. Thus Λ is aperiodic as claimed.

Since Λ is aperiodic there exist $\mu, \nu \in \Lambda$ such that $s_{\mu^*}as_\nu = tp_w$ for some $w \in \Lambda^0$ and some $t \in K \setminus \{0\}$ (see Lemma 2.3(2)). Since w connects to a line point, there exists $\gamma \in w\Lambda P_l(\Lambda)$. Thus $p_{s(\gamma)} \in I_{\overline{P_l(\Lambda)}}$ and then $s_\nu s_\gamma = s_\nu s_\gamma p_{s(\gamma)} \in I_{\overline{P_l(\Lambda)}}$. Therefore

$$0 \neq p_{s(\gamma)} = s_{\gamma^*}s_\gamma = s_{\gamma^*}p_ws_\gamma = t^{-1}s_{\gamma^*}s_{\mu^*}as_\nu s_\gamma \in t^{-1}s_{\gamma^*}s_{\mu^*}aI_{\overline{P_l(\Lambda)}}.$$

Thus $aI_{\overline{P_l(\Lambda)}} \neq \{0\}$ as well.

Conversely, suppose that $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = I_{\overline{P_l(\Lambda)}}$ is essential. Pick $w \in \Lambda^0$. Since $I_{\overline{P_l(\Lambda)}}$ is essential, there exists $b \in I_{\overline{P_l(\Lambda)}}$ such that $p_wb \neq 0$. By [10, Lemma 5.4], $I_{\overline{P_l(\Lambda)}} = \operatorname{span}\{s_\alpha s_{\beta^*} : s(\alpha) = s(\beta) \in \overline{P_l(\Lambda)}\}$ so we can write $b = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta}s_\alpha s_{\beta^*}$ with $s(\alpha) = s(\beta) \in \overline{P_l(\Lambda)}$. Since $p_wb \neq 0$, there exists $(\mu,\nu) \in F$ such that $p_ws_\mu s_{\nu^*} \neq 0$. For this μ we have $\mu \in w\Lambda\overline{P_l(\Lambda)}$. Since every element in $\overline{P_l(\Lambda)}$ connects to $P_l(\Lambda)$ by Lemma 2.5, we can pick $\eta \in s(\mu)\Lambda P_l(\Lambda)$. Thus $\mu\eta$ connects w to $P_l(\Lambda)$. □

The next corollary follows immediately from Theorem 3.10(1) and the lattice isomorphism $H \mapsto I_H$ of the saturated hereditary subsets of Λ^0 onto the graded ideals of $\mathrm{KP}_K(\Lambda)$ from [10, Theorem 5.1].

Corollary 3.11. *Let Λ be a row-finite k-graph with no sources and let K be a field.*

(1) $\operatorname{Soc}(\mathrm{KP}_K(\Lambda))$ *is a graded ideal of* $\mathrm{KP}_K(\Lambda)$.
(2) $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = \{0\}$ *if and only if* $P_l(\Lambda) = \emptyset$.
(3) $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = \mathrm{KP}_K(\Lambda)$ *if and only if* $\overline{P_l(\Lambda)} = \Lambda^0$.

Examples 3.12. (1) Let Λ be a row-finite k-graph with one vertex v. Then $P_l(\Lambda) = \emptyset$ and hence $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = \{0\}$ (for if there is just one infinite path then it must be periodic). In particular, the Kumjian-Pask algebras of the 2-graphs with one vertex, N_1 blue edges and N_2 red edges considered by Davidson and Yang in [17] all have zero socle.

(2) Every vertex of Ω_k is a line point, and hence $\operatorname{Soc}(\mathrm{KP}_K(\Omega_k)) = \mathrm{KP}_K(\Omega_k)$.

4. SEMISIMPLE KUMJIAN-PASK ALGEBRAS

Throughout this section Λ is a row-finite k-graph with no sources and K is a field. A semiprime ring R is *semisimple* if $\operatorname{Soc}(R) = R$. By [13, Remark 4.3.5], $\operatorname{Soc}(R)$ is a direct sum of minimal left-ideals. Thus R is semisimple if and only if it is a direct sum of its minimal left-ideals. By Corollary 3.11, $\mathrm{KP}_K(\Lambda)$ is semisimple if and only if the saturated hereditary closure of the line points is all of Λ^0. In this section we study semisimple Kumjian-Pask algebras. The proof of the next lemma is very similar to the proof of its Leavitt path algebra version [4, Lemma 2.2].

Lemma 4.1. *Let e be an idempotent in $\mathrm{KP}_K(\Lambda)$. Then the corner $e\,\mathrm{KP}_K(\Lambda)e$ is a division ring if and only if it is isomorphic to K as a ring.*

Proof. Suppose $e\,\mathrm{KP}_K(\Lambda)e$ is a division ring. By Lemma 2.7, $\mathrm{KP}_K(\Lambda)$ is semiprime. Now [13, Proposition 4.3.3] applies, giving that $\mathrm{KP}_K(\Lambda)e$ is a minimal left-ideal. By Theorem 3.9(2), there exists $v \in P_l(\Lambda)$ such that $\mathrm{KP}_K(\Lambda)e$ is isomorphic to $\mathrm{KP}_K(\Lambda)p_v$ as a left $\mathrm{KP}_K(\Lambda)$-module. Lemma 3.8 says that $p_v\,\mathrm{KP}_K(\Lambda)p_v$ is isomorphic to Kp_v as a ring; since Kp_v and K are isomorphic, so are $p_v\,\mathrm{KP}_K(\Lambda)p_v$ and K.

For an idempotent f in a ring R, the map $\phi : fRf \to \mathrm{End}_R(Rf)$, where $\phi(frf)(tf) = tfrf$, is a ring isomorphism. Thus

$$e\,\mathrm{KP}_K(\Lambda)e \cong \mathrm{End}_{\mathrm{KP}_K(\Lambda)}(\mathrm{KP}_K(\Lambda)e) \cong \mathrm{End}_{\mathrm{KP}_K(\Lambda)}(\mathrm{KP}_K(\Lambda)p_v) \cong p_v\,\mathrm{KP}_K(\Lambda)p_v \cong K.$$

The converse is obvious. □

To state our main theorem we need a few definitions. Suppose V is a cancellative abelian monoid. An element $u \in V$ is a *unit* if there exists $v \in V$ such that $u + v = 0$. An element $s \in V$ is an *atom* (or irreducible) if $s = u + v$ implies u or v is a unit; V is *atomic* if every element of the monoid is a sum of atoms, and V satisfies the *refinement property* if $a + b = c + d$ in V implies there exist $z_1, z_2, z_3, z_4 \in V$ such that $a = z_1 + z_2$, $b = z_3 + z_4$, $c = z_1 + z_3$ and $d = z_2 + z_4$.

For a ring R, denote by $V(R)$ the monoid of finitely generated projective modules over R. The direct sum of nonzero projective modules is nonzero, so the only unit in $V(R)$ is 0. Thus $s \in V(R)$ is an atom if $s = u + v$ implies that either u or v is zero. Let e and f be idempotents in R. Since $R \cong Re \oplus R(1-e)$, Re is a projective module. We say e and f are *equivalent* if $Re \cong Rf$, and we denote the equivalence class of e in R by $[e]$. If there exist $a, b \in R$ such that $e = ab$ and $f = ba$ then the map $Re \to Rf : re \mapsto rea$ is an isomorphism of R-modules with inverse $rf \mapsto rfb$, and hence e and f are equivalent. We identify the set of equivalence classes of idempotents in R with a submonoid of $V(R)$ via $[e] \mapsto Re$.

Let R be a ring with local units. There is an order on idempotents in R where $e \leq f$ if and only if $ef = e$. We claim that $e \leq f$ if and only if $Re \subset Rf$. If $e \leq f$ then $aef = ae$ for all $a \in R$, so $Re \subset Rf$. Conversely, if $Re \subset Rf$ then $e \in Rf$ since R has local units. So $e = af$ for some $a \in R$ and hence $ef = aff = af = e$. Thus $e \leq f$, completing the claim.

A ring R is *von Neumann regular* if for every $a \in R$ there exists a $b \in R$ such that $aba = a$. We need the following lemma to prove Theorem 4.3.

Lemma 4.2. *If $V(\mathrm{KP}_K(\Lambda))$ is an atomic monoid with the refinement property then for every $y \in \Lambda^\infty$ there exists $m \in \mathbb{N}^k$ such that $y(m)\Lambda^\infty = \{\sigma^m(y)\}$.*

Proof. By way of contradiction, suppose there exists $y \in \Lambda^\infty$ such that $|y(m)\Lambda^\infty| > 1$ for all $m \in \mathbb{N}^k$. Let $v = y(0)$. Using Remark 3.3(1), there exists a sequence $\{m_i\} \subset \mathbb{N}^k$ such that $m_{i+1} \geq m_i$ and $|y(m_i)\Lambda^{m_{i+1}-m_i}| > 1$. For $i \in \mathbb{N} \setminus \{0\}$ set

$$e_i := s_{y(0,m_i)} s_{y(0,m_i)^*}.$$

We claim $\{p_v\,\mathrm{KP}_K(\Lambda)e_i\}$ is a strictly decreasing sequence of left ideals in $p_v\,\mathrm{KP}_K(\Lambda)p_v$.

Let $i \in \mathbb{N} \setminus \{0\}$. To see $p_v \operatorname{KP}_K(\Lambda)e_i$ contains $p_v \operatorname{KP}_K(\Lambda)e_{i+1}$ note that

$$\begin{aligned} e_{i+1}e_i &= s_{y(0,m_{i+1})}s_{y(0,m_{i+1})^*}s_{y(0,m_i)}s_{y(0,m_i)^*} \\ &= s_{y(0,m_{i+1})}s_{y(m_i,m_{i+1})^*}s_{y(0,m_i)^*}s_{y(0,m_i)}s_{y(0,m_i)^*} \\ &= s_{y(0,m_{i+1})}s_{y(m_i,m_{i+1})^*}s_{y(0,m_i)^*} \quad \text{(using (KP3))} \\ &= s_{y(0,m_{i+1})}s_{y(0,m_{i+1})^*} = e_{i+1} \quad \text{(using (KP2))}. \end{aligned}$$

This shows that $e_i \geq e_{i+1}$ and that $p_v \operatorname{KP}_K(\Lambda)e_i \supset p_v \operatorname{KP}_K(\Lambda)e_{i+1}$. To see the containment is strict we will show that $e_i \notin p_v \operatorname{KP}_K(\Lambda)e_{i+1}$. Since $|y(m_i)\Lambda^{m_{i+1}-m_i}| > 1$, we can pick $\mu_i \in y(m_i)\Lambda^{m_{i+1}-m_i} \setminus \{y(m_i, m_{i+1})\}$. Consider $\nu_i = y(0, m_i)\mu_i$. Then, using (KP3),

$$e_i s_{\nu_i} = s_{y(0,m_i)}s_{y(0,m_i)^*}s_{\nu_i} = s_{y(0,m_i)}s_{\mu_i} = s_{\nu_i} \neq 0.$$

But for any $a \in \operatorname{KP}_K(\Lambda)$

$$p_v a e_{i+1} s_{\nu_i} = p_v a s_{y(0,m_{i+1})}s_{y(m_i,m_{i+1})^*}s_{y(0,m_i)^*}s_{y(0,m_i)}s_{\mu_i} = 0$$

by (KP3) because μ_i has degree $m_{i+1} - m_i$. Thus $p_v a e_{i+1} \neq e_i$ for all $a \in \operatorname{KP}_K(\Lambda)$ and hence $e_i \notin p_v \operatorname{KP}_K(\Lambda)e_{i+1}$. Therefore $p_v \operatorname{KP}_K(\Lambda)e_i \supsetneq p_v \operatorname{KP}_K(\Lambda)e_{i+1}$ and $e_i \gneq e_{i+1}$. Now e_{i+1} and $e_i - e_{i+1}$ are nonzero orthogonal idempotents and, denoting an orthogonal sum with $\oplus$, we have $e_i = e_{i+1} \oplus (e_i - e_{i+1})$.

Since $V(\operatorname{KP}_K(\Lambda))$ is atomic by assumption, there exists N and atoms $q_1, \dots, q_N \in V(\operatorname{KP}_K(\Lambda))$ so that $[e_1] = q_1 + \cdots + q_N$. For this N, we have

$$e_1 = e_{N+1} \oplus (e_N - e_{N+1}) \oplus \cdots \oplus (e_1 - e_2).$$

So in $V(\operatorname{KP}_K(\Lambda))$ we have

$$[e_{N+1}] + [e_N - e_{N+1}] + \cdots + [e_1 - e_2] = q_1 + \cdots + q_N.$$

We note here for future use that $[e_{N+1}] \neq 0$ because $e_{N+1} = s_{y(0,m_{N+1})}s_{y(0,m_{N+1})^*}$ is equivalent to $s_{y(0,m_{N+1})^*}s_{y(0,m_{N+1})} = p_{y(m_{N+1})} \neq 0$.

By assumption $V(\operatorname{KP}_K(\Lambda))$ has the refinement property, so there exist $z_{j,l}$, $1 \leq j \leq N, 1 \leq l \leq N+1$ in $V(\operatorname{KP}_K(\Lambda))$ such that

$$q_j = \sum_l z_{j,l}, \quad [e_l - e_{l+1}] = \sum_j z_{j,l} \text{ for } 1 \leq l \leq N \quad \text{and} \quad [e_{N+1}] = \sum_j z_{j,N+1}.$$

Since q_j is an atom for each j, there exists a unique $l_j \in \{1, \dots N+1\}$ such that $z_{j,l_j} \neq 0$. In other words, there are precisely N nonzero $z_{j,l}$. So there exists an $l_0 \in \{1, \dots, N+1\}$ such that $z_{j,l_0} = 0$ for all j. But $[e_{N+1}] \neq 0$, so $l_0 \neq N+1$ and $[e_{l_0} - e_{l_0+1}] = 0$. Now $\{0\} = \operatorname{KP}_K(\Lambda) = \operatorname{KP}_K(\Lambda)(e_{l_0} - e_{l_0+1})$. By nondegeneracy of $\operatorname{KP}_K(\Lambda)$ we have $e_{l_0} = e_{l_0+1}$, a contradiction. Hence there exists m such that $|y(m)\Lambda^\infty| = 1$. □

In [4] Abrams, Aranda Pino, Perera and Siles Molina "characterize the semisimple Leavitt path algebras by describing them in categorical, ring-theoretic, graph-theoretic, and explicit terms." They start by proving in [4, Theorem 2.3] that a semiprime ring R with local units is semisimple if and only if eight equivalent conditions hold. Among these conditions are that R is locally left (respectively, right) artinian, that every corner of R is left (respectively, right) artinian, and that R is isomorphic to a certain direct sum of matrix algebras over corners of R. In [4, Theorem 2.4] they restrict their attention to the Leavitt path algebra $L(E)$ of a directed graph E and link the conditions of [4, Theorem 2.3] to $L(E)$ being categorically left (respectively, right) artinian, $L(E)$ being von Neumann regular and properties of the monoid $V(L(E))$, and purely graph-theoretic conditions. Our Theorem 4.3 is the higher-rank analogue of [4, Theorem 2.4], but we have omitted some of the many equivalent conditions of [4, Theorem 2.4] to concentrate on those pertinent to $\mathrm{KP}_K(\Lambda)$ and Λ in particular.

Theorem 4.3. *Let Λ be a row-finite k-graph with no sources and K be a field. The following conditions are equivalent.*

1. $\mathrm{KP}_K(\Lambda)$ *is semisimple.*
2. *There exists a countable set Υ and for each $i \in \Upsilon$ there exists $n_i \in \mathbb{N} \cup \{\infty\}$ such that $\mathrm{KP}_K(\Lambda) \cong \bigoplus_{i\in\Upsilon} M_{n_i}(K)$.*
3. *There exists a countable set Υ and for each $i \in \Upsilon$ there exists $n_i \in \mathbb{N} \cup \{\infty\}$ such that $V(\mathrm{KP}_K(\Lambda)) \cong \oplus_{i\in\Upsilon}\mathbb{N}$, and $\mathrm{KP}_K(\Lambda)$ is von Neumann regular.*
4. *$V(\mathrm{KP}_K(\Lambda))$ is cancellative and atomic, and $\mathrm{KP}_K(\Lambda)$ is von Neumann regular.*
5. *For every $y \in \Lambda^\infty$, there exists $m \in \mathbb{N}^k$ such that $y(m) \in P_l(\Lambda)$.*

Remark 4.4. Condition (5) of Theorem 4.3 implies that every infinite path is aperiodic. To see this, suppose $y \in \Lambda^\infty$ and pick m such that $y(m) \in P_l(\Lambda)$. Then $y(m)\Lambda^\infty = \{\sigma^m(y)\}$ with $\sigma^m(y)$ aperiodic. Since $\sigma^m(y)$ is aperiodic, so is y.

Proof of Theorem 4.3. We will show (1) ⇒ (2) ⇒ (3) ⇒ (4) ⇒ (5) ⇒ (1).

(1) ⇒ (2). Recall that an idempotent e in a semiprime ring R is minimal if Re is a minimal left-ideal [13, p. 143]. Theorem 2.3 of [4] says that a semiprime ring R is semisimple if and only if there exists an index set Υ and for each $i \in \Upsilon$ there exist minimal idempotents $e_i \in R$ and $n_i \in \mathbb{N}\cup\{\infty\}$ such that $R \cong \sum_{i\in\Upsilon} M_{n_i}(e_iRe_i)$. Since Re_i is a minimal left-ideal in R, [13, Proposition 4.3.3] shows that e_iRe_i is a division ring. Assume $\mathrm{KP}_K(\Lambda)$ is semisimple. By Lemma 2.7, $\mathrm{KP}_K(R)$ is semiprime. Applying [4, Theorem 2.3] to $R = \mathrm{KP}_K(\Lambda)$ we obtain Υ, e_i and n_i as above. By Lemma 2.4, $\mathrm{KP}_K(\Lambda)$ has a countable set of local units, and thus Υ is countable by [4, Theorem 2.3]. By Lemma 4.1, each $e_i \,\mathrm{KP}_K(\Lambda)e_i$ is isomorphic to K. This gives (2).

(2) ⇒ (3). Assume (2). Each $M_{n_i}(K)$ is von Neumann regular, and hence so is the direct sum. Since $V(M_{n_i}(K)) = \mathbb{N}$ and $V(R_1 \oplus R_2) = V(R_1) \oplus V(R_2)$, (3) holds.

(3) ⇒ (4). This follows because $\oplus_{i\in\Upsilon}\mathbb{N}$ cancellative and atomic.

(4) ⇒ (5). By [8, 3rd example on p. 412], "π-regular rings" are exchange rings. Since π-regular rings are a generalization of von Neumann regular rings it follows that von Neumann regular rings are exchange rings. By [6, Proposition 1.2], if R is an exchange ring then $V(R)$ has the refinement property.

Assume (4). Then $V(\mathrm{KP}_K(\Lambda))$ has the refinement property. Let $y \in \Lambda^\infty$. By Lemma 4.2, there exists $t \in \mathbb{N}^k$ such that $y(t)\Lambda^\infty = \{\sigma^t(y)\}$. We claim that $y(t)$ is a line point.

By way of contradiction, suppose that $y(t)$ is not a line point. Since $y(t)\Lambda^\infty = \{\sigma^t(y)\}$, $\sigma^t(y)$ must be periodic. For convenience set $z := y(t,\infty)$; then $z(0)\Lambda^\infty = \{z\}$ and z is periodic. So there exists $m \neq n \in \mathbb{N}^k$ such that $z(m) = z(n)$.

Let $\pi_h : \mathbb{N}^k \to \mathbb{N}$ be the projection onto the h-th factor. By switching m and n if necessary, we may assume that there exists an $l \in \{1,\dots,k\}$ such that $\pi_l(m-n) > 0$. Let $v = z(0)$. Consider the element $p_v + s_{z(0,m)}s_{z(0,n)^*}$. Since we assumed that $\mathrm{KP}_K(\Lambda)$ is von Neumann regular, there exists $b \in \mathrm{KP}_K(\Lambda)$ such that

$$(p_v + s_{z(0,m)}s_{z(0,n)^*})b(p_v + s_{z(0,m)}s_{z(0,n)^*}) = p_v + s_{z(0,m)}s_{z(0,n)^*}. \tag{4.1}$$

Since both sides of (4.1) are nonzero we can assume $b \in p_v \,\mathrm{KP}_K(\Lambda)p_v$. Write b in normal form as $b = \sum_{(\alpha,\beta)\in F} r_{\alpha,\beta}s_\alpha s_{\beta^*}$. Then for all $(\alpha,\beta),(\mu,\nu) \in F$ we have $d(\beta) = d(\nu)$, and, since $|v\Lambda^\infty| = 1$, we have $\beta = \nu$. So we can assume that

$$b = \sum_{i=1}^{N} r_i s_{\alpha_i} s_{\beta^*} \quad r_i \in K \setminus \{0\}$$

for some fixed β. Furthermore, we can assume that the α_i are ordered so that $\pi_l(d(\alpha_i)) \leq \pi_l(d(\alpha_{i'}))$ for all $1 \leq i \leq i' \leq N$.

We will obtain a contradiction by comparing the degrees on the left and right sides of (4.1). Define the π_l-degree of a homogeneous element to be the l-th component of its degree. The minimal π_l-degree associated to the left side of (4.1) is obtained by the monomial $p_v s_{\alpha_1} s_{\beta^*} p_v = s_{\alpha_1} s_{\beta^*}$. The minimal π_l-degree appearing in the right side of (4.1) is zero. So $\pi_l(d(\alpha_1) - d(\beta)) = 0$. The maximal π_l-degree on the left hand side of (4.1) is obtained by the monomial $s_{z(0,m)} s_{z(0,n)^*} s_{\alpha_N} s_{\beta^*} s_{z(0,m)} s_{z(0,n)^*}$ whose π_l-degree is $2\pi_l(m-n) + \pi_l(d(\alpha_N) - d(\beta))$. By Lemma 3.6, $s_{z(0,m)} s_{z(0,n)^*} s_{\alpha_N} s_{\beta^*} s_{z(0,m)} s_{z(0,n)^*} \neq 0$. Since the maximal π_l-degree on the right hand side of (4.1) is $\pi_l(m-n) > 0$, we have $2\pi_l(m-n) + \pi_l(d(\alpha_N) - d(\beta)) = \pi_l(m-n)$. Thus $\pi_l(d(\alpha_N) - d(\beta)) = -\pi_l(m-n) < 0$. Now $0 = \pi_l(d(\alpha_1) - d(\beta)) \leq \pi_l(d(\alpha_N) - d(\beta)) < 0$, a contradiction. Thus $z = \sigma^t(y)$ is aperiodic, and $y(t)$ is a line point as claimed. This gives (5).

(5) $\Rightarrow$ (1). Assume (5). By Corollary 3.11, we need to show that $\overline{P_l(\Lambda)} = \Lambda^0$. By way of contradiction suppose there exists $v \in \Lambda^0 \setminus \overline{P_l(\Lambda)}$. We define a path $y \in v\Lambda^\infty$ by recursion. Since v is in the complement of the saturated set $\overline{P_l(\Lambda)}$, there exists a path $\mu_1 \in v\Lambda^{(1,\dots,1)}$ such that $s(\mu_1) \notin \overline{P_l(\Lambda)}$. Suppose that μ_i has been defined is such a way that $r(\mu_i) = s(\mu_{i-1})$, $d(\mu_i) = (1,\dots,1)$ and $s(\mu_i) \notin \overline{P_l(\Lambda)}$. Because $\overline{P_l(\Lambda)}$ is saturated, there exists $\mu_{i+1} \in s(\mu_i)\Lambda^{(1,\dots,1)}$ such that $s(\mu_{i+1}) \notin \overline{P_l(\Lambda)}$. Set $y = \mu_1\mu_2 \cdots \in v\Lambda^\infty$. By (5), there exists $m \in \mathbb{N}^k$ such that $y(m) \in P_l(\Lambda)$. But for i sufficiently large, $i \cdot (1,\dots,1) > m$ and since $P_l(\Lambda)$ is hereditary, $s(\mu_i) \in P_l(\Lambda)$, a contradiction. Thus $\Lambda^0 = \overline{P_l(\Lambda)}$ and hence $\mathrm{KP}_K(\Lambda)$ is semisimple. □

Remark 4.5. A simple Kumjian-Pask algebra need not be semisimple. To see this, pick any row-finite, aperiodic and cofinal k-graph Λ which contains a periodic path. Then $\mathrm{KP}_K(\Lambda)$ is simple by [10, Theorem 6.1], but is not semisimple by Theorem 4.3.

Our next goal is to find a more precise description of the matrix decomposition of semisimple Kumjian-Pask algebras given by item (2) of Theorem 4.3. We will show that $\mathrm{KP}_K(\Lambda)$ is isomorphic to $\bigoplus_{[v]\in P_l(\Lambda)/\sim} I_{\overline{\{v\}}}$ where $\sim$ is an equivalence relation on the set of line points and $I_{\overline{\{v\}}}$ is the graded ideal associated to the saturated hereditary subset $\overline{\{v\}}$ of Λ^0.

Let $v, w \in P_l(\Lambda)$ so that $v\Lambda^\infty = \{x\}$ and $w\Lambda^\infty = \{y\}$. Define a relation on $P_l(\Lambda)$ by

$$v \sim w \Leftrightarrow \exists\ m, n \in \mathbb{N}^k \text{ such that } x(m) = y(n).$$

It follows from item (2) of Lemma 4.6 that $\sim$ defines an equivalence relation on $P_l(\Lambda)$.

Let H be a saturated hereditary subset of Λ^0 and recall that the *quotient graph* $\Lambda \setminus H$ is

$$\Lambda \setminus H := (\Lambda^0 \setminus H, s^{-1}(\Lambda^0 \setminus H), r|, s|); \tag{4.2}$$

$\Lambda \setminus H$ is a row-finite k-graph with no sources by [10, Lemma 5.3].

Lemma 4.6. *Let $v, w \in P_l(\Lambda)$, say $v\Lambda^\infty = \{y\}$ and $w\Lambda^\infty = \{z\}$.*

(1) If $u \in \overline{\{v\}}$, then $\overline{\{u\}} = \overline{\{v\}}$.
(2) $\overline{\{w\}} \cap \overline{\{v\}} \neq \emptyset \quad \Leftrightarrow \quad \overline{\{w\}} = \overline{\{v\}} \quad \Leftrightarrow \quad w \sim v$.

(3) If Λ *contains no periodic paths then the ideal* $I_{\overline{\{v\}}}$ *is minimal.*

Proof. (1) Suppose $u \in \overline{\{v\}}$. Then $\overline{\{u\}} \subset \overline{\{v\}}$. For the reverse inclusion it suffices to show that $v \in \overline{\{u\}}$. Recall that $\Sigma^0(\{v\})$ is the hereditary closure of $\{v\}$. Since $u \in \overline{\{v\}}$, by Lemma 2.5, there exists $\mu \in u\Lambda\Sigma^0(\{v\})$. But $\overline{\{u\}}$ is hereditary, so $s(\mu) \in \overline{\{u\}} \cap \Sigma^0(\{v\})$. Because $\Sigma^0(\{v\})$ is the hereditary closure of $\{v\}$ there exists a path $\lambda \in v\Lambda s(\mu) \subset v\Lambda\overline{\{u\}}$. Since $v \in P_l(\Lambda)$, $|v\Lambda^{d(\lambda)}| = 1$ by Remark 3.3(1). That is, $v\Lambda^{d(\lambda)} = \{\lambda\}$. Hence $s(v\Lambda^{d(\lambda)}) \subset \overline{\{u\}}$ which implies $v \in \overline{\{u\}}$ because $\overline{\{u\}}$ is saturated.

(2) First, assume $\overline{\{w\}} \cap \overline{\{v\}} \neq \emptyset$. Let $u \in \overline{\{w\}} \cap \overline{\{v\}}$. By (1), we have $\overline{\{w\}} = \overline{\{u\}} = \overline{\{v\}}$.

Second, assume $\overline{\{w\}} = \overline{\{v\}}$. Then $v \in \overline{\{w\}}$, so there exists $\mu \in v\Lambda\Sigma^0(\{w\})$ by Lemma 2.5. Since $s(\mu) \in \Sigma^0(\{w\})$, there exists $\nu \in w\Lambda s(\mu)$. Now $v\Lambda^\infty = \{y\}$ and $w\Lambda^\infty = \{z\}$ so $\mu = y(0, d(\mu))$ and $\nu = z(0, d(\nu))$. But $y(d(\mu)) = s(\mu) = s(\nu) = z(d(\nu))$, and hence $v \sim w$.

Third, assume $v \sim w$. There exist $n, m \in \mathbb{N}^k$ such that $y(m) = z(n)$. But $\overline{\{v\}}$ and $\overline{\{w\}}$ are hereditary, so $y(m) \in \overline{\{v\}} \cap \overline{\{w\}} \neq \emptyset$.

(3) Now suppose that Λ contains no periodic paths. Then $\Lambda \setminus H$ is aperiodic for all saturated hereditary subsets H of Λ^0, and [10, Theorem 5.6] implies that every ideal of $\mathrm{KP}_K(\Lambda)$ is graded. Thus $H \mapsto I_H$ is an isomorphism of the lattice of saturated hereditary subsets of Λ onto the lattice of ideals of $\mathrm{KP}_K(\Lambda)$. By (1), $\overline{\{v\}}$ is a minimal saturated hereditary subset, and hence the ideal $I_{\overline{\{v\}}}$ is a minimal ideal. □

Next we want to show that the ideals $\{I_{\overline{\{v\}}}\}_{[v] \in P_l(\Lambda)/\sim}$ are mutually orthogonal. For ideals I, J in a ring R we let $IJ := \{ab : a \in I, b \in J\}$ and $I \cdot J := \operatorname{span}\{ab : a \in I, b \in J\}$. Then $I \cdot J$ is a two-sided ideal in R, and $IJ \subset I \cdot J \subset I \cap J$.

Lemma 4.7. *Let* H *and* L *be saturated hereditary subsets of* Λ^0. *Then* $I_H \cap I_L = I_{H \cap L} = I_H \cdot I_L$. *In particular, if* $H \cap L = \emptyset$ *then* $I_H I_L = \{0\}$.

Proof. The second statement follows immediately from the first because $I_H I_L \subset I_H \cdot I_L = I_{H \cap L}$. Recall that $H \mapsto I_H$ is an isomorphism of the lattice of saturated hereditary subsets of Λ onto the lattice of graded ideals of $\mathrm{KP}_K(\Lambda)$. Since the intersection of graded ideals is graded, $I_H \cap I_L = I_W$ for some saturated hereditary subset W of Λ^0. Using that I_W is generated by p_v with $v \in W$, it is straightforward to show that $W = H \cap L$. Thus $I_H \cap I_L = I_{H \cap L}$

By definition of $I_H \cdot I_L$ we have $I_H \cdot I_L \subset I_H \cap I_L = I_{H \cap L}$. For the reverse inclusion, let $p_v \in I_{H \cap L}$. Then $v \in H \cap L$, and hence $p_v \in I_H$ and $p_v \in I_L$. Now $p_v = p_v^2 \in I_H \cdot I_L$. Since $I_{H \cap L}$ is generated by p_v with $v \in H \cap L$ and $I_H \cdot I_L$ is an ideal we get $I_{H \cap L} \subset I_H \cdot I_L$. □

We now have another corollary of Theorem 3.10.

Corollary 4.8. *Let* Λ *be a row-finite* k*-graph with no sources and let* K *be a field. If* $\mathrm{KP}_K(\Lambda)$ *semisimple then* $\mathrm{KP}_K(\Lambda) = \bigoplus_{[v] \in P_l(\Lambda)/\sim} I_{\overline{\{v\}}}$.

Proof. Let $v, w \in P_l(\Lambda)$ such that $v \not\sim w$. By Lemma 4.6(2), $\overline{\{v\}} \cap \overline{\{w\}} = \emptyset$, so by Lemma 4.7 we have $I_{\overline{\{v\}}} I_{\overline{\{w\}}} = \{0\}$. Thus the internal direct sum $\bigoplus_{[v] \in P_l(\Lambda)/\sim} I_{\overline{\{v\}}}$ is an ideal in $\mathrm{KP}_K(\Lambda)$.

Since $\mathrm{KP}_K(\Lambda)$ is semisimple, $\operatorname{Soc}(\mathrm{KP}_K(\Lambda)) = \mathrm{KP}_K(\Lambda)$. But $\operatorname{Soc}(\mathrm{KP}_K(\Lambda))$ is the ideal generated by $\{p_v : v \in P_l(\Lambda)\}$ by Theorem 3.10. Thus $\mathrm{KP}_K(\Lambda) \subset \bigoplus_{[v] \in P_l(\Lambda)/\sim} I_{\overline{\{v\}}}$, that is, $\mathrm{KP}_K(\Lambda) = \bigoplus_{[v] \in P_l(\Lambda)/\sim} I_{\overline{\{v\}}}$. □

Next we analyze the ideals $I_{\overline{\{v\}}}$ for $v \in P_l(\Lambda)$.

Lemma 4.9. *(1) If $v \in P_l(\Lambda)$ then $I_{\overline{\{v\}}}$ contains a subalgebra isomorphic to $M_\infty(K)$.*
(2) If $\mathrm{KP}_K(\Lambda)$ is both simple and semisimple then $\mathrm{KP}_K(\Lambda) \cong M_\infty(K)$.
(3) If $\mathrm{KP}_K(\Lambda)$ is semisimple then $I_{\overline{\{v\}}} \cong M_\infty(K)$ for all $v \in P_l(\Lambda)$.

Proof. (1) Let $v \in P_l(\Lambda)$. Then $v\Lambda^\infty = \{y\}$. It suffices to show $I_{\overline{\{v\}}}$ contains a set of matrix units indexed by $\mathbb{N} \times \mathbb{N}$. For $i, j \in \mathbb{N}$ define

$$e_{i,j} = \begin{cases} s_{y((i,0,\dots,0),(j,0,\dots,0))} & \text{if } i < j \\ s_{y((j,0,\dots,0),(i,0,\dots,0))^*} & \text{if } j < i \\ p_{y(j,0,\dots,0)} & \text{if } i = j. \end{cases}$$

We claim $\{e_{i,j}\}$ forms a set of matrix units in $I_{\overline{\{v\}}}$. First note that $e_{i,j}e_{h,\ell} = 0$ unless $j = h$. A case by case analysis shows $e_{i,j}e_{j,\ell} = e_{i,\ell}$. For example: suppose $i < \ell < j$. Then $|v\Lambda^{j-\ell}| = 1$ since $v \in P_l(\Lambda)$. Thus

$$\begin{aligned} e_{i,j}e_{j,\ell} &= s_{y((i,0,\dots,0),(j,0,\dots,0))} s_{y((\ell,0,\dots,0),(j,0,\dots,0))^*} \\ &= s_{y((i,0,\dots,0),(\ell,0,\dots,0))} s_{y((\ell,0,\dots,0),(j,0,\dots,0))} s_{y((\ell,0,\dots,0),(j,0,\dots,0))^*} \end{aligned}$$

which by Remark 3.3(4) is

$$= s_{y((i,0,\dots,0),(\ell,0,\dots,0))} p_{y(\ell,0,\dots,0)} = e_{i,\ell}.$$

The other cases follow from similar arguments. Thus $\{e_{i,j}\}_{i,j\in\mathbb{N}}$ forms a set of matrix units in $I_{\overline{\{v\}}}$, and hence $M_\infty(K)$ is isomorphic to a subalgebra of $I_{\overline{\{v\}}}$.

(2) Suppose that $\mathrm{KP}_K(\Lambda)$ is semisimple. Then $\mathrm{KP}_K(\Lambda) \cong \bigoplus_{i\in\Upsilon} M_{n_i}(K)$ by Theorem 4.3((1) $\Rightarrow$ (2)). Since $\mathrm{KP}_K(\Lambda)$ is simple this direct sum has only one summand $M_m(K)$ for some $m \in \mathbb{N} \cup \{\infty\}$. By (1), $M_\infty(K)$ is a subalgebra of $\mathrm{KP}_K(\Lambda)$. Since $M_\infty(K)$ cannot be isomorphic to a subalgebra of $M_n(K)$ for $n \in \mathbb{N}$, $m = \infty$ as desired.

(3) Suppose that $\mathrm{KP}_K(\Lambda)$ is semisimple. We will show that $I_{\overline{\{v\}}}$ is isomorphic to a simple Kumjian-Pask algebra and then invoke (2) to get the result. Since $\mathrm{KP}_K(\Lambda)$ is semisimple it follows from Corollary 4.8 that $I_{\overline{\{v\}}} \cong \mathrm{KP}_K(\Lambda)/J$ where

$$J = \bigoplus_{[w]\in(P_l(\Lambda)/\sim)\setminus\{[v]\}} I_{\overline{\{w\}}}.$$

It follows from Theorem 4.3(5) that Λ has no periodic paths (see Remark 4.4). So for any saturated hereditary set H, $\Lambda \setminus H$ is aperiodic, and therefore every ideal of $\mathrm{KP}_K(\Lambda)$ is graded by [10, Theorem 5.6]. In particular J is graded, and hence $J = I_L$ for some saturated hereditary subset L of Λ^0 by [10, Theorem 5.1]. Now by [10, Proposition 5.5], $I_{\overline{\{v\}}} \cong \mathrm{KP}_K(\Lambda \setminus L)$.

Next we will show that $\mathrm{KP}_K(\Lambda \setminus L)$ is simple and semisimple. Since $\mathrm{KP}_K(\Lambda)$ is semisimple every infinite path $y \in \Lambda^\infty$ contains a line point by Theorem 4.3. But now every infinite path $y \in (\Lambda \setminus L)^\infty$ contains a line point as well, so $\mathrm{KP}_K(\Lambda \setminus L)$ is semisimple by Theorem 4.3.

As observed above, $\Lambda \setminus L$ is aperiodic. Theorem 6.1 of [10] says that $\mathrm{KP}_K(\Lambda \setminus L)$ is simple if and only if $\Lambda \setminus L$ is cofinal. Pick $x \in (\Lambda \setminus L)^\infty$, $w \in (\Lambda \setminus L)^0$ and $y \in w(\Lambda \setminus L)^\infty$. We need to show that there exists a t such that $y(0, t)$ connects w to x.

Since we know that x, y both contain line points, there exists $m, n \in \mathbb{N}^k$ such that $x(m), y(n) \in P_l(\Lambda)$. Since $x(m), y(n) \notin L$ we have $p_{x(m)}, p_{y(n)} \notin J$, and hence $x(m), y(n) \in \overline{\{v\}}$. Thus $x(m) \sim y(n)$ by Lemma 4.6(2). By definition of $\sim$ there exist

m', n' such that $x(m+m') = \sigma^m(x)(m') = \sigma^n(y)(n') = y(n+n')$. Now $y(0, n+n')$ connects w to x and hence $\Lambda \setminus L$ is cofinal. Thus $\mathrm{KP}_K(\Lambda \setminus L)$ is simple. Now we apply (2) to see $I_{\overline{\{v\}}} \cong \mathrm{KP}_K(\Lambda \setminus L) \cong M_\infty(K)$. □

The next result sharpens condition (2) of Theorem 4.3.

Theorem 4.10. *Let Λ be a row-finite k-graph with no sources and K be a field. Then $\mathrm{KP}_K(\Lambda)$ is semisimple if and only if $\mathrm{KP}_K(\Lambda) \cong \bigoplus_{[v]\in(P_l(\Lambda)/\sim)} M_\infty(K)$.*

Proof. Suppose that $\mathrm{KP}_K(\Lambda)$ is semisimple. Then Corollary 4.8 gives that $\mathrm{KP}_K(\Lambda) = \bigoplus_{[v]\in(P_l(\Lambda)/\sim)} I_{\overline{\{v\}}}$ and so $\mathrm{KP}_K(\Lambda) \cong \bigoplus_{[v]\in(P_l(\Lambda)/\sim)} M_\infty(K)$ by Lemma 4.9(3). The converse is immediate from Theorem 4.3 ((2) $\Rightarrow$ (1)). □

Remark 4.11. Theorem 4.10 shows that every semisimple Kumjian-Pask algebra can be obtained as a Leavitt path algebra, that is, the added generality of k-graphs gives nothing new for semisimple algebras. For $n \in \mathbb{N} \cup \{\infty\}$ it is easy to construct a 1-graph Λ_n whose Kumjian-Pask algebra is isomorphic to $\bigoplus_{i=1}^n M_\infty(K)$. Just take Λ_n to be the following graph with n vertical components.



By inspection $\overline{P_l(\Lambda_n)} = \Lambda_n^0$ and $v \sim w \in P_l(\Lambda)$ if and only if v and w are in the same vertical component. Thus by Theorem 4.10, $\mathrm{KP}_K(\Lambda_n) \cong \bigoplus_{i=1}^n M_\infty(K)$.

References

[1] Gene Abrams, Gonzalo Aranda Pino, and Mercedes Siles Molina, *Finite-dimensional Leavitt path algebras*, J. Pure Appl. Algebra **209** (2007), no. 3, 753–762.

[2] Gene Abrams and Gonzalo Aranda Pino, *The Leavitt path algebra of a graph*, J. Algebra **293** (2005), no. 2, 319–334.

[3] Gene Abrams and Gonzalo Aranda Pino, *Purely infinite simple Leavitt path algebras*, J. Pure Appl. Algebra **207** (2006), no. 3, 553–563.

[4] Gene Abrams, Gonzalo Aranda Pino, Francesc Perera, and Mercedes Siles Molina, *Chain conditions for Leavitt path algebras*, Forum Math. **22** (2010), no. 1, 95–114.

[5] Gene Abrams, Kulumani M. Rangaswamy, and Mercedes Siles Molina, *The socle series of a Leavitt path algebra*, Israel J. Math. **184** (2011), 413–435.

[6] Pere Ara, Kenneth. R. Goodearl, Kevin C. O'Meara, and Enrique Pardo, *Separative cancellation for projective modules over exchange rings*, Israel J. Math. **105** (1998), 105–137.

[7] Pere Ara, Maria A. Moreno, and Enrique Pardo, *Nonstable K-theory for graph algebras*, Algebr. Represent. Theory **10** (2007), no. 2, 157–178.

[8] Pere Ara, *Extensions of exchange rings*, J. Algebra **197** (1997), no. 2, 409–423.

[9] Pere Ara, Martin Mathieu, and Eduard Ortega, *The maximal C^*-algebra of quotients as an operator bimodule*, Arch. Math. (Basel) **92** (2009), no. 5, 405–413.

[10] Gonzalo Aranda Pino, John Clark, Astrid an Huef, and Iain Raeburn, *Kumjian-Pask algebras of higher-rank graphs*, to appear in Trans. Amer. Math. Soc., arXiv:1106.4361v1.

[11] Gonzalo Aranda Pino, Dolores Martín Barquero, Cándido Martín González, and Mercedes Siles Molina, *Socle theory for Leavitt path algebras of arbitrary graphs*, Rev. Mat. Iberoam. **26** (2010), no. 2, 611–638.

[12] Gonzalo Aranda Pino, Dolores Martín Barquero, Cándido Martín González, and Mercedes Siles Molina, *The socle of a Leavitt path algebra*, J. Pure Appl. Algebra **212** (2008), no. 3, 500–509.

[13] Konstantin I. Beidar, Wallace S. Martindale III, and Aleksandr V. Mikhalev, *Rings with generalized identities*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 196, Marcel Dekker Inc., New York, 1996.

[14] Paul E. Bland, *Rings and their modules*, Walter de Gruyter & Co., Berlin, 2011.

[15] Bernhard Burgstaller, *The K-theory of certain C^*-algebras endowed with gauge actions*, J. Funct. Anal. **256** (2009), no. 6, 1693–1707.

[16] Kenneth R. Davidson, Stephen C. Power, and Dilian Yang, *Atomic representations of rank 2 graph algebras*, J. Funct. Anal. **255** (2008), no. 4, 819–853.

[17] Kenneth R. Davidson and Dilian Yang, *Periodicity in rank 2 graph algebras*, Canad. J. Math. **61** (2009), no. 6, 1239–1261.

[18] D. Gwion Evans and Aidan Sims, *When is the Cuntz-Krieger algebra of a higher-rank graph approximately finite-dimensional?*, arXiv:1112.4549v1.

[19] Cynthia Farthing, Paul S. Muhly, Trent Yeend *Higher-rank graph C^*-algebras: an inverse semigroup and groupoid approach*, Semigroup Forum **71** (2005), no. 2, 159–187.

[20] Robert Hazlewood, Iain Raeburn, Aidan Sims and Samuel B.G. Webster, *On some fundamental results about higher-rank graphs and their C^*-algebras*, arXiv:1110.2269v1.

[21] David W. Kribs and Stephen C. Power, *The analytic algebras of higher rank graphs*, Math. Proc. R. Ir. Acad. **106A** (2006), no. 2, 199–218.

[22] Alex Kumjian and David Pask, *Higher rank graph C^*-algebras*, New York J. Math. **6** (2000), 1–20.

[23] Alex Kumjian, David Pask, Iain Raeburn, and Jean Renault, *Graphs, groupoids, and Cuntz-Krieger algebras*, J. Funct. Anal. **144** (1997), no. 2, 505–541.

[24] William G. Leavitt, *Modules without invariant basis number*, Proc. Amer. Math. Soc. **8** (1957), 322–328.

[25] David Pask, Iain Raeburn, Mikael Rørdam, and Aidan Sims, *Rank-two graphs whose C^*-algebras are direct limits of circle algebras*, J. Funct. Anal. **239** (2006), no. 1, 137–178.

[26] Iain Raeburn, Aidan Sims, and Trent Yeend, *Higher-rank graphs and their C^*-algebras*, Proc. Edinb. Math. Soc. (2) **46** (2003), no. 1, 99–115.

[27] David I. Robertson and Aidan Sims, *Simplicity of C^*-algebras associated to row-finite locally convex higher-rank graphs*, Israel J. Math. **172** (2009), 171–192.

[28] Guyan Robertson and Tim Steger, *Affine buildings, tiling systems and higher rank Cuntz-Krieger algebras*, J. Reine Angew. Math. **513** (1999), 115–144.

[29] Mercedes Siles Molina, *Algebras of quotients of path algebras*, J. Algebra **319** (2008), 5265–5278.

[30] Adam Skalski and Joachim Zacharias, *Entropy of shifts on higher-rank graph C^*-algebras*, Houston J. Math. **34** (2008), no. 1, 269–282.

[31] Aidan Sims, Gauge-invariant ideals in the C^*-algebras of finitely aligned higher-rank graphs, Canad. J. Math. **58** (2006), no. 6, 1268–1290.

[32] Mark Tomforde, *Uniqueness theorems and ideal structure for Leavitt path algebras*, J. Algebra **318** (2007), no. 1, 270–299.

[33] Trent Yeend, *Groupoid models for the C^*-algebras of topological higher-rank graphs*, J. Operator Theory **57** (2007), no. 1, 95–120,

Department of Mathematics and Statistics, University of Otago, Dunedin 9054, New Zealand

E-mail address: `jbrown@maths.otago.ac.nz, astrid@maths.otago.ac.nz`